

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/48, 1/42, G01N 33/68, 33/53	A1	(11) International Publication Number: WO 99/29894 (43) International Publication Date: 17 June 1999 (17.06.99)
---	----	---

(21) International Application Number: PCT/US98/25196
(22) International Filing Date: 2 December 1998 (02.12.98)
(30) Priority Data:
60/067,833 5 December 1997 (05.12.97) US
(71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).
(72) Inventors; and
(75) Inventors/Applicants (for US only): EPPS, Dennis, E. [US/US]; 240 Point 'O' Woods Drive, Portage, MI 49002 (US). MARSCHKE, Charles, K. [US/US]; 3210 Olney Street, Kalamazoo, MI 49006 (US).
(74) Agent: KERBER, Lori, L.; Pharmacia & Upjohn Company, Intellectual Property Legal Services, 301 Henrietta Street, Kalamazoo, MI 49001 (US).


(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: FLUORESCENCE-BASED HIGH THROUGHPUT SCREENING ASSAYS FOR PROTEIN KINASES AND PHOSPHATASES



(57) Abstract

The invention relates to novel fluorescence-based assays for protein kinases and phosphatases which can be used in high throughput screening. The methods of the invention utilize a competitive immunoassay to determine the amount of substrate that is phosphorylated or dephosphorylated during the course of a kinase or phosphatase reaction to yield a product, as well as the phosphorylating or dephosphorylating activity of a kinase or phosphatase.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Fluorescence-Based High Throughput Screening Assays for Protein Kinases and Phosphatases

FIELD OF THE INVENTION

5

The invention relates to novel fluorescence-based assays for kinases and phosphatases which can be used in high throughput screening.

BACKGROUND OF THE INVENTION

10

Eukaryotes employ phosphorylation and dephosphorylation of specific proteins to regulate many cellular processes (T. Hunter, *Cell* 80:225-236 (1995); (Karin, M., *Curr. Opin. Cell Biol.* 3: 467-473 (1991)). These processes include signal transduction, cell division, and initiation of gene transcription. Thus, significant events in an organism's maintenance, adaptation, and susceptibility to disease are controlled by protein phosphorylation and dephosphorylation. These phenomena are so extensive that it has been estimated that humans have around 2,000 protein kinase genes and 1,000 protein phosphatase genes (T. Hunter, *Cell* 80:225-236 (1995)), some of these likely coding for disease susceptibility. For these reasons, protein kinases and phosphatases are good targets for the development of drug therapies.

The most frequently used protein kinase and phosphatase screens employ either radioactive ATP or ELISAs. However, the use of radioactive ATP is undesirable due to the attendant costs of record-keeping, waste-disposal, and the fact that the assay format is not homogeneous. ELISAs are undesirable because they have a lower assay throughput due to the extra steps required for both washing and the enzyme reaction.

Fluorescence detection in the visible wavelengths offer an alternative to the use of radiotracers or ELISAs for kinase and phosphatase assays, as fluorescence offers detection limits comparable to those of radioactivity. Furthermore, this eliminates the cost of radioactive waste disposal. For example, the change in absorbance and fluorescence spectra of phosphotyrosine which occurs upon dephosphorylation has been used for the continuous monitoring of protein-tyrosine phosphatase (PTP) activity (Zhao, Z. *et al.*, *Anal. Biochem.* 202:361-366 (1993)). However, previously developed fluorometric assays for kinases and phosphatases have not been especially amenable to the requirements of high throughput screening.

Fluorescence detection frequently offers the advantage of using homogeneous assay formats (i.e. - "mix, incubate, and read"). Indeed, the high throughput screening

(HTS) field is moving rapidly toward the use of fluorescence, luminescence, absorbance, and other optical methods. Two fluorescence techniques, fluorescence polarization (FP) and fluorescence resonance energy transfer (FRET) are finding widespread use for assays, both in the private sector for HTS, secondary assays including kinetics, SAR studies, *etc.*, and in university laboratories. The use of FP is particularly desirable since its readout is independent of the emission intensity (Checovich, W.J., *et al.*, *Nature* 375:254-256 (1995); Dandliker, W.B., *et al.*, *Methods in Enzymology* 74:3-28 (1981)) and is thus insensitive to the presence of colored compounds that quench fluorescence emission. FRET, although susceptible to quenching, can also be used effectively, especially for continuous enzyme assays.

From the forgoing, it will be clear that there is a continuing need for the development of cost-effective, facile, and sensitive optical kinase and phosphatase assays for both high throughput screening (HTS) and secondary assays.

15 INFORMATION DISCLOSURE

Checovich, W.J., *et al.*, *Nature* 375:254-256 (1995).

Dandliker, W.B., *et al.*, *Methods in Enzymology* 74:3-28 (1981).

E. Harlow and D. Lane, eds., *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory (1988).

T. Hunter, *Cell* 80:225-236 (1995).

Leavine, L.M., *et al.*, *Anal. Biochem.* 247:83-88 (1997).

Owicki, J.C., *Genetic Engineering News* 17:27 (November 1, 1997).

Rotman, B., *et al.*, *Proc. Nat. Acad. Sci.* 50:1-6 (1963).

Seethala, R. and R. Menzel, *A Fluorescence Polarization Tyrosine Kinase Assay for High Throughput Screening*, 3rd Annual Conference of The Society for Biomolecular Screening, San Diego, CA, September 22-25, 1997.

SUMMARY OF THE INVENTION

The invention relates to novel fluorescence-based assays for protein kinases and phosphatases which can be used in high throughput screening. The methods of the invention utilize a competitive immunoassay to determine the amount of substrate that is phosphorylated or dephosphorylated during the course of a kinase or phosphatase
5 reaction to yield a product, as well as the phosphorylating or dephosphorylating activity of a kinase or phosphatase.

Thus, in one embodiment, the invention relates to a method of determining the phosphorylating activity of an enzyme comprising the steps of:

- (a) combining the enzyme with
 - 10 (i) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein the amino acid is selected from the group consisting of serine, threonine and tyrosine;
 - (ii) a substrate molecule comprising the same amino acid that is phosphorylated in said reporter, wherein said substrate molecule is capable of being
15 phosphorylated at said amino acid by said enzyme to yield a product;
 - (iii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid; and
 - (iv) a high-energy phosphate source;
- (b) measuring the fluorescence polarization (FP), FQ, or fluorescence resonance
20 spectroscopy (FCS) of the reporter following the combination of step (a); and
- (c) using the FP, FQ, or FCS measurement of step (b) to determine the activity of the enzyme.

In another embodiment, the invention relates to a method for determining the dephosphorylating activity of an enzyme comprising the steps of:

- 25 (a) combining the enzyme with
 - (i) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein the amino acid is selected from the group consisting of serine, threonine and tyrosine;
 - (ii) a substrate molecule comprising the same phosphorylated amino acid
30 as said reporter, wherein said substrate molecule is capable of being dephosphorylated at said amino acid by said enzyme to yield a product; and
 - (iii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid;
- (b) measuring the FP, FQ, or FCS of said reporter following the combination of
35 step (a); and
- (c) using the FP, FQ, or FCS measurement of step (b) to determine the activity

of the enzyme.

The methods of the invention can also be used to determine the phosphorylation or dephosphorylation of a substrate molecule by an enzyme. Thus, in another embodiment, the invention relates to a method for determining the phosphorylation of
5 a substrate molecule by an enzyme at an amino acid selected from the group consisting of serine, threonine and tyrosine, comprising the steps of:

- (a) combining the substrate molecule with
 - (i) the enzyme
 - (ii) a reporter molecule comprising a fluorescent label and a
10 phosphorylated amino acid, wherein the amino acid is the same amino acid which is phosphorylated in the reporter;
 - (iii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid; and
 - (iv) a high-energy phosphate source;
- 15 (b) measuring the FP, FQ, or FCS of the reporter following the combination of step (a); and
- (c) using the FP, FQ, or FCS measurement of step (b) to determine whether the substrate molecule has been phosphorylated.

In another embodiment, the invention relates to a method for determining the
20 dephosphorylation of a substrate molecule by an enzyme, wherein the substrate molecule comprises a phosphorylated amino acid, and wherein the amino acid is selected from the group consisting of serine, threonine and tyrosine, comprising the steps of:

- (a) combining the substrate molecule with
25 (i) the enzyme;
- (ii) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein the reporter molecule comprises the same phosphorylated amino acid as the substrate molecule; and
- (iii) an antibody which selectively binds to a molecule comprising the
30 phosphorylated amino acid;
- (b) measuring the FP, FQ, or FCS of the reporter following the combination of step (a); and
- (c) using the FP, FQ, or FCS measurement of step (b) to determine whether the substrate molecule has been dephosphorylated.

35 In a preferred embodiment, the substrate in any of the above methods is combined with the enzyme before the addition of the reporter and the antibody. In

another preferred embodiment, the substrate, the reporter, and the antibody are combined with the enzyme simultaneously.

Because the above-described methods of the invention utilize a competitive immunoassay to determine the amount of phosphorylated or dephosphorylated substrate (*i.e.*, the amount of product) produced, the amount of phosphorylated substrate required to displace the reporter from the antibody will vary depending upon the K_d of the phosphorylated substrate for the antibody and the K_d of the antibody for the reporter molecule.

Thus, where the K_d of the phosphorylated substrate for the antibody is, *e.g.*, 10-fold higher than the K_d of the antibody for the reporter molecule, then an amount of phosphorylated substrate ten times higher than the amount of reporter will be required for the phosphorylated substrate to displace the reporter from the antibody.

In a more preferred embodiment, the K_d of the phosphorylated substrate for the antibody will be approximately equal to the K_d of the antibody for the reporter molecule. In a still more preferred embodiment, the K_d of the phosphorylated substrate for the antibody will be less than the K_d of the antibody for the reporter molecule. In this situation, phosphorylation of the substrate will quantitatively displace the reporter from the antibody.

In accordance with the above description, one way of reducing the amount of substrate needed to displace the reporter from the antibody is to choose a reporter having a low K_d for the antibody. Because anti-phosphorylamino acid antibodies may have a higher affinity for a fluorescently labeled phosphorylamino acid than for a fluorescently labeled peptide comprising the same phosphorylamino acid, in a preferred embodiment, such peptides are used as the reporter.

The methods of the invention also allow the utilization of a continuous recording assay (*i.e.*, a "real time" assay) for the determination of either kinase or phosphatase activity by using a FRET format.

Thus, in another embodiment, the invention relates to a method of determining the phosphorylating activity of an enzyme comprising the steps of:

(a) combining the enzyme with:

(i) a substrate molecule comprising an amino acid selected from the group consisting of Ser, Thr, and Tyr, wherein said substrate molecule is capable of being phosphorylated at said amino acid by said enzyme to yield a product, and wherein said substrate molecule is labeled with an acceptor fluorophore;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with a donor fluorophore which

corresponds to the acceptor fluorophore labeling said substrate; and

(iii) a high-energy phosphate source;

(b) measuring the FRET of the combination of step (a); and

(c) using the FRET measurement of step (b) to determine the activity of the
5 enzyme.

In another embodiment, the invention relates to a method of determining the phosphorylating activity of an enzyme comprising the steps of:

(a) combining the enzyme with:

(i) a substrate molecule comprising an amino acid selected from the
10 group consisting of Ser, Thr, and Tyr, wherein said substrate molecule is capable of being phosphorylated at said amino acid by said enzyme to yield a product, and wherein said substrate molecule is labeled with a donor fluorophore;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore
15 which corresponds to the donor fluorophore labeling said substrate; and

(iii) a high-energy phosphate source;

(b) measuring the FRET of the combination of step (a); and

(c) using the FRET measurement of step (b) to determine the activity of the
enzyme.

20 In another embodiment, the invention relates to a method of determining the dephosphorylating activity of an enzyme comprising the steps of:

(a) combining the enzyme with:

(i) a substrate molecule comprising a phosphorylated amino acid selected from the group consisting of phosphoserine, phosphothreonine and phosphotyrosine,
25 wherein said substrate molecule is labeled with an acceptor fluorophore;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with a donor fluorophore which corresponds to the acceptor fluorophore labeling said substrate; and

(iii) a high-energy phosphate source;

30 (b) measuring the FRET of the combination of step (a); and

(c) using the FRET measurement of step (b) to determine the activity of the
enzyme.

In another embodiment, the invention relates to a method of determining the dephosphorylating activity of an enzyme comprising the steps of:

35 (a) combining the enzyme with:

(i) a substrate molecule comprising a phosphorylated amino acid selected

from the group consisting of phosphoserine, phosphothreonine and phosphotyrosine, wherein said substrate molecule is labeled with a donor fluorophore;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore
5 which corresponds to the donor fluorophore labeling said substrate; and

(iii) a high-energy phosphate source;

(b) measuring the FRET of the combination of step (a); and

(c) using the FRET measurement of step (b) to determine the activity of the enzyme.

10 In another embodiment, the invention relates to a method for determining the phosphorylation of a substrate molecule by an enzyme at an amino acid selected from the group consisting of serine, threonine and tyrosine, wherein said substrate molecule is labeled with an acceptor fluorophore, comprising the steps of:

(a) combining the substrate molecule with

15 (i) the enzyme

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with a donor fluorophore which corresponds to the acceptor fluorophore labeling said substrate; and

(iii) a high-energy phosphate source;

20 (b) measuring the FRET of the reporter following the combination of step (a);
and

(c) using the FRET measurement of step (b) to determine whether the substrate molecule has been phosphorylated.

In another embodiment, the invention relates to a method for determining the
25 phosphorylation of a substrate molecule by an enzyme at an amino acid selected from the group consisting of serine, threonine and tyrosine, wherein said substrate molecule is labeled with a donor fluorophore, comprising the steps of:

(a) combining the substrate molecule with:

(i) the enzyme

30 (ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore which corresponds to the donor fluorophore labeling said substrate; and

(iii) a high-energy phosphate source;

(b) measuring the FRET of the report r following the combination of step (a);
35 and

(c) using the FRET measurement of step (b) to determine whether the substrate

molecule has been phosphorylated.

In another embodiment, the invention relates to a method for determining the dephosphorylation of a substrate molecule by an enzyme, wherein the substrate molecule comprises a phosphorylated amino acid selected from the group consisting of phosphoserine, phosphothreonine and phosphotyrosine, and wherein said substrate
5 molecule is labeled with an acceptor fluorophore comprising the steps of:

(a) combining the substrate molecule with:

(i) the enzyme;

(ii) an antibody which selectively binds to a molecule comprising the
10 phosphorylated amino acid, said antibody being labeled with a donor fluorophore which corresponds to the acceptor fluorophore labeling said substrate;

(b) measuring the FRET of the reporter following the combination of step (a);

and

(c) using the FRET measurement of step (b) to determine whether the substrate
15 molecule has been dephosphorylated.

In another embodiment, the invention relates to a method for determining the dephosphorylation of a substrate molecule by an enzyme, wherein the substrate molecule comprises a phosphorylated amino acid selected from the group consisting of serine, threonine and tyrosine, and wherein said substrate molecule is labeled with a
20 donor fluorophore comprising the steps of:

(a) combining the substrate molecule with:

(i) the enzyme;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore
25 which corresponds to the donor fluorophore labeling said substrate;

(b) measuring the FRET of the reporter following the combination of step (a);

and

(c) using the FRET measurement of step (b) to determine whether the substrate molecule has been dephosphorylated.

30 The methods of the invention can also be used to identify an agent capable of increasing or decreasing the phosphorylating activity of an enzyme comprising the steps of:

(a) performing the above method of determining the phosphorylating activity of an enzyme in the presence and in the absence of the agent;

35 (b) comparing the activity of the enzyme in the presence of the agent with the activity of the enzyme in the absence of the agent to determine whether the

phosphorylating activity of the enzyme in the presence of the agent is increased or decreased.

In yet another embodiment, the invention relates to a method of screening for an agent capable of increasing or decreasing the dephosphorylating activity of an enzyme comprising the steps of:

- (a) performing the above method of determining the dephosphorylating activity of an enzyme in the presence and in the absence of said agent;
- (b) comparing the activity of said enzyme in the presence of said agent with the activity of said enzyme in the absence of said agent to determine whether the dephosphorylating activity of said enzyme in the presence of said agent is increased or decreased.

BRIEF DESCRIPTION OF THE FIGURES

15

Figures 1A-1G show the chemical structures of reagents used in Example 1.

Figures 2A-2C: Figures 2A-2C are graphs showing the binding of sigma antibodies to fluoresceinated phosphorylamino acids. Fixed concentrations of P-Tyr-F (25.8 nM)(Fig. 2A), P-Ser-F (5.9 nM)(Fig. 2B), or P-Thr-F (10 nM)(Fig. 2C) were titrated with the specific antibodies as described in Methods of Example 1. The polarization was recorded after each addition and the data analyzed using Equation 7 of Example 1 in conjunction with a nonlinear least squares fitting program. In this and subsequent figures, the solid lines represent the theoretical fits to the experimental data.

Figures 3A and 3B: Figures 3A and 3B are graphs showing the binding of MBL anti-phosphotyrosine antibody to P-Tyr-F. A fixed concentration (15 nM) of P-Tyr-F was titrated with the MBL antibody as described in Methods of Example 1. The FP (Fig. 3A) and total polarized emission (Fig. 3B) were monitored during the titration. The quenching data were analyzed by nonlinear least squares fitting using Equation 7 of Example 1, and the FP data using Equation 20 of Example 1 with the values of C and Q_c/Q_L calculated from Equation 7 of Example 1 substituted into it.

Figures 4A-4C: Figures 4A-4C are graphs showing the displacement of fluoresceinated phosphorylamino acids from Sigma antibodies by the corresponding unlabelled phosphorylamino acids. Concentrations of the fluoresceinated phosphorylamino acids and the corresponding Sigma antibodies were fixed, and their displacement by the phosphorylamino acids was followed by measuring the decrease in the fluorescence polarization. The fixed concentrations were: P-Tyr-F, 1 nM and MAB,

2.05 nM sites (Fig. 4A); P-Ser-F, 1 nM and MAb, 1 nM sites (Fig. 4B); P-Thr-F, 150 nM and MAb, 150 nM sites (Fig. 4C).

Figures 5A and 5B: Figures 5A and 5B are graphs showing the competitive displacement of P-Tyr-F from MBL antibody by phosphoryltyrosine as measured by FQ and FP. MBL antibody, 7.1 nM sites and P-Tyr-F, 1.7 nM were preincubated, and displacement initiated by the addition of a concentrated solution of phosphotyrosine. The increase in emission intensity (Fig. 5A) and decrease in FP (Fig. 5B) were monitored as a function of added phosphoryltyrosine. The data were analyzed as described in Figure 3.

Figure 6 is a graph showing the competitive displacement of P-Tyr-F from Sigma antibody by phosphorylated JAK-2 kinase peptides. Sigma antibody, 2.05 nM sites and P-Tyr-F, 1 nM, were preincubated and displacement initiated as described for Figure 5. The polarization data were analyzed as described in Figure 3. ●--●, JAK-2(a); ○--○, JAK-2(b).

Figures 7A-7D: Figures 7A-7D are graphs showing the competitive displacement of P-Tyr-F from MBL antibody by phosphorylated JAK-2 peptide substrates JAK-2(a)(Figs. 7A and 7B) and JAK-2(b)(Figs. 7C and 7D). All procedures were as described in Figure 5. For both peptides, fixed antibody = 7.1 nM sites and P-Tyr-F = 1.7 nM.


DETAILED DESCRIPTION OF THE INVENTION

The invention relates to novel fluorescence-based assays for kinases and phosphatases which can be used in high throughput screening. As used herein, the term "kinase" refers to an enzyme capable of phosphorylating its substrate at a Ser, Thr, or Tyr residue, while the term "phosphatase" refers to an enzyme capable of dephosphorylating its substrate at a phosphoserine, phosphothreonine, or phosphotyrosine residue. The methods of the invention utilize a competitive immunoassay to determine the amount of phosphorylated or dephosphorylated substrate produced during the course of a kinase or phosphatase reaction, as well as the phosphorylating or dephosphorylating activity of a kinase or phosphatase. Unless otherwise indicated, "phosphorylating activity" as used herein is synonymous with "kinase activity," and "dephosphorylating activity" as used herein is synonymous with "phosphatase activity." Similarly, unless otherwise indicated, a "kinase" is defined herein as a biological material capable of phosphorylating a peptide or protein, and a "phosphatase" is defined her in as a biological material capable of dephosphorylating a peptide or protein. Further, where the enzyme used in any of the assays of the

invention is a kinase, the term "phosphorylated substrate" is synonymous with "product," (*i.e.*, the product derived from the enzymatic reaction). Similarly, where the enzyme used in any of the assays of the invention is a phosphatase, the term "dephosphorylated substrate" is synonymous with "product," (*i.e.*, the product derived from the enzymatic reaction).

In the methods of the invention, a reporter molecule comprising a fluorescent label and a phosphorylamino acid (P-AA) selected from the group consisting of Ser, Thr and Tyr (hereinafter referred to as a "reporter molecule," or "reporter") competes with a phosphorylated substrate molecule, comprising the same P-AA as the reporter molecule, for an antibody specific for the P-AA. The antibody and reporter molecule are chosen so that binding of the antibody to the reporter causes a change in the reporter which is detectable using FP, FQ, or FCS. Knowledge of the concentration of reporter and substrate used, the dissociation constant (K_d) of the phosphorylated substrate for the antibody, the K_d of the reporter for the antibody, and the change in the fluorescent properties of the reporter will allow calculation of the amount of phosphorylated substrate present, and the determination of kinase or phosphatase activity, as is described below.

Thus, by the methods of the invention, phosphorylation of a substrate peptide or protein by a kinase can be monitored by specific displacement of a reporter molecule from an antibody by the reaction product of the kinase assay (the phosphorylated substrate molecule). One assay format for the fluorescent kinase assay is given below.



In this new scheme, a kinase reaction is carried out in a reaction mixture by contacting a kinase with a substrate molecule in the presence of a high energy

phosphate source such as ATP or GTP. The reaction is ~~allowed to~~ proceed for a period of time, and is stopped (for example, by the addition of a metal chelator such as EDTA or EGTA). Subsequently, both antibody and reporter are added to the reaction mixture, whereby the reaction product (the phosphorylated substrate molecule) specifically
5 displaces the reporter molecule from the antibody.

In another embodiment, the antibody and reporter molecule are present at time = 0, so that the phosphorylated substrate competes with the reporter for the antibody, giving intermediate values of polarization and/or quenching, thus providing a homogeneous format for high-throughput screening. Of course, one of ordinary skill
10 will realize that the antibody and reporter molecule can only be present at the time that the enzyme is added where neither the antibody nor the reporter are a substrate for the enzyme.

Thus, the invention provides a method of determining the phosphorylating activity of an enzyme comprising the steps of:

- 15 (a) combining said enzyme with
- (i) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein said amino acid is selected from the group consisting of serine, threonine and tyrosine;
 - (ii) a substrate capable of being phosphorylated by said enzyme at the
20 same amino acid which is phosphorylated in said reporter;
 - (iii) an anti-phosphorylamino acid antibody which is specific for said phosphorylated amino acid, and which selectively binds to a molecule comprising said phosphorylated amino acid; and
 - (iv) a high energy phosphate source
- 25 (b) measuring the FP, FQ, or FCS of said reporter following the combination of step (a); and
- (c) using the FP, FQ, or FCS measurement of step (b) to determine the activity of said enzyme.

In one embodiment, the substrate is combined with the enzyme before the
30 addition of said reporter and said antibody. In another embodiment, the substrate, the reporter, and the antibody are combined with the enzyme simultaneously.

As described, the dephosphorylation of a substrate peptide or protein can be also be measured by the competitive immunoassays of the present invention. Thus, in another embodiment, the invention provides a method for determining the
35 dephosphorylating activity of an enzyme comprising the steps of:

- (a) combining said enzyme with

(i) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein said amino acid is selected from the group consisting of serine, threonine and tyrosine;

(ii) a substrate comprising the same phosphorylated amino acid as said reporter, wherein said substrate is capable of being dephosphorylated at said amino acid by said enzyme; and

(iii) an antibody which selectively binds to a molecule comprising said phosphorylated amino acid;

(b) measuring the FP, FQ, or FCS of said reporter following the combination of step (a); and

(c) using the FP, FQ, or FCS measurement of step (b) to determine the dephosphorylating activity of the enzyme.

As in the kinase assay described above, the antibody and reporter molecule may be added after the phosphatase reaction has proceeded for some time, in which case the remaining phosphorylated substrate will specifically displace the reporter molecule from the antibody. Alternatively, the antibody and reporter molecule may be present at time = 0, so that the phosphorylated substrate competes with the reporter for the antibody, giving intermediate values of polarization and/or quenching. As is true for the kinase assay, the antibody and reporter molecule can only be present at the time that the enzyme is added where neither the antibody nor the reporter are a substrate for the enzyme.

Because the methods of the invention utilize a competitive immunoassay to determine the amount of phosphorylated or dephosphorylated substrate (i.e., the amount of product) produced, the amount of phosphorylated substrate required to displace the reporter from the antibody will vary depending upon the K_d of the phosphorylated substrate for the antibody and the K_d of the antibody for the reporter molecule. Thus, where the K_d of the phosphorylated substrate for the antibody is, e.g., 10-fold less than the K_d of the antibody for the reporter molecule, then an amount of phosphorylated substrate ten times higher than the amount of reporter will be required for the phosphorylated substrate to displace the reporter from the antibody. Where the K_d of the phosphorylated substrate for the antibody is approximately equal to the K_d of the antibody for the reporter molecule, only 50% of the maximal signal can be achieved. In a highly preferred embodiment, the K_d of the phosphorylated substrate for the antibody will be less than the K_d of the antibody for the reporter molecule. In this situation, phosphorylation of the substrate will quantitatively displace the reporter from the antibody.

Another way of reducing the amount of substrate relates to the choice of reporter molecule used in the methods of the invention. A suitable reporter molecule to be used in the methods of the invention is a fluorescently-labeled molecule, preferably a peptide or protein, comprising a phosphorylated amino acid, wherein said amino acid is selected from the group consisting of serine, threonine, and tyrosine. Suitable reporter molecules for use in the methods of the present invention thus include fluorescently labeled phosphoserine, phosphothreonine or phosphotyrosine, as well as an appropriately fluorescently labeled peptide comprising a phosphorylamino acid (P-AA) selected from the group consisting of Ser, Thr and Tyr. Selection of reporter wherein the K_d of the antibody for the reporter is higher than the K_d of the phosphorylated substrate for the antibody will reduce the amount of substrate needed to displace the reporter from the antibody. Because anti-phosphorylamino acid antibodies may have a higher affinity for a fluorescently labeled phosphorylamino acid than for a fluorescently labeled peptide comprising the same phosphorylamino acid, in a preferred embodiment, such peptides are used as the reporter.

Production of peptides to be used as reporters may be accomplished by any one of a number of methods that are well known to those of ordinary skill, such as by enzymatic cleavage, chemical synthesis, or expression of a recombinantly produced peptide. The reporter peptide may be synthesized and then phosphorylated, or instead the phosphorylated amino acid or amino acids may be incorporated into the peptide at the time that the peptide is synthesized. Phosphorylamino acids for incorporation into chemically synthesized peptides may be obtained from numerous commercial sources, such as Sigma (St. Louis, MO). In preferred embodiment, labeling of the reporter is accomplished by including a Cys residue in the sequence one to two residues away from the phosphorylatable amino acid.

Suitable fluorescent labels to be used in the methods of the invention include any fluorophore that, upon the binding of the reporter molecule by an anti-phosphorylamino acid antibody, undergoes a change detectable by FP, FQ, or FCS. Of course, in order to be used in the methods of the invention, the label cannot interfere with recognition of the reporter by the anti-phosphorylamino acid antibody. Where the detection method to be used is FP, the fluorescent label to be used may be any probe which, when combined with a molecule comprising a phosphorylated amino acid (P-AA) selected from the group consisting of Ser, Thr and Tyr to form a reporter molecule, undergoes a change in fluorescence lifetime when the reporter molecule binds to a larger molecule (i.e., the antibody). Where the detection method to be used is FQ, the fluorescent label to be used may be any environment-sensitive probe which, when

combined with a molecule comprising a phosphorylamino acid (P-AA) selected from the group consisting of Ser, Thr and Tyr to form a reporter molecule, undergoes a change in fluorescence intensity when the reporter molecule binds to a larger molecule (*i.e.*, the antibody). In FCS, the difference in the diffusion coefficients of two bound molecules
5 (with the smaller of the two being fluorescently labeled) is observed in a very small volume. Thus, the smaller, labeled molecule will diffuse into the observed volume faster in the unbound state than it will if bound to a larger molecule. For this technique, any probe that undergoes minimal photobleaching can be used, with preference given to those with the highest quantum yields.

10 Utilization of a FRET assay format results in a continuous recording assay for either phosphatase or kinase activities. Where a FRET format is to be used, the antibody is labeled with a suitable donor or acceptor fluorophore, while the substrate is labeled with the donor or acceptor fluorophore that complements the fluorophore labeling the antibody. For example, the substrate peptide can be labeled with a donor
15 fluorophore such as fluorescein or Oregon green, while the antibody is labeled with a suitable fluorophore acceptor, such as rhodamine, with excitation at the absorbance maximum of the donor, and emission observed at the maximum of either fluorophore. Thus, as the target amino acid of the labeled substrate becomes phosphorylated, the labeled antibody will bind to the substrate, resulting in both the quenching of the donor
20 fluorescence and the enhancement of the acceptor fluorescence.

An antibody suitable for use in the FP, FQ, FRET and FCS methods of the invention is one that binds specifically to phosphoserine, phosphothreonine, or phosphotyrosine, and that produces changes in either the intrinsic polarization or quenching of the emission intensity of the fluoresceinated version of the
25 phosphorylamino acid to which it binds. Furthermore, where the assay format used is FP, FQ, or FCS, the difference between the K_d of the antibody for the phosphorylated substrate and the K_d of the antibody for the reporter molecule will dictate the amount of phosphorylated substrate that must be present in order for the phosphorylated substrate to displace the reporter from the antibody. Thus, where the K_d of the
30 phosphorylated substrate for the antibody is, *e.g.*, 10-fold less than the K_d of the antibody for the reporter molecule, then an amount of phosphorylated substrate ten times higher than the amount of reporter will be required for the phosphorylated substrate to displace the reporter from the antibody. Where the K_d of the phosphorylated substrate for the antibody is approximately equal to the K_d of the
35 antibody for the reporter molecule, only 50% of the maximal signal can be achieved. In a highly preferred embodiment, the K_d of the phosphorylated substrate for the

antibody will be less than the K_d of the antibody for the reporter molecule. In this situation, phosphorylation of the substrate will quantitatively displace the reporter from the antibody.

Because the methods of the invention are to be used to assay kinase and
5 phosphatase activity, it is preferred that displacement of the reporter molecule from the antibody be accomplished with a relatively low concentration of phosphorylated substrate. As is described above, this may be accomplished by using a reporter having a K_d for the antibody that is higher than the K_d of the phosphorylated substrate for the antibody, and will allow detection of lower concentrations of phosphorylated substrate.
10 With proper selection of the reporter, the assay format can be FP, FQ, or FCS.

Anti-phosphorylamino acid antibodies may be obtained from numerous commercial sources, including Sigma (St. Louis, MO), ICN Biomedicals (Costa Mesa, CA), Life Technologies (Gaithersburg, MD), Transduction Labs, (Lexington, KY), Molecular Biology Laboratories (Nagoya, Japan), Upstate Biologicals (Lake Placid, NY)
15 and Zymed Laboratories (South San Francisco, CA). In addition, anti-phosphorylamino acid antibodies may be prepared as described in Zoppini *et al.* (*Eur. J. Lab. Med.* 1(2): 101-103 (1993)), and in the references cited therein. Determination of whether a certain anti-P-AA antibody produces changes in either the intrinsic polarization or quenching of the emission intensity of the F-P-AA may be made according to the
20 method described in Example 1.

As used herein, the term "antibody" includes monoclonal antibodies, polyclonal antibodies, single chain antibodies, and ligand-binding fragments of antibodies, such as Fab and F(ab')₂. Various procedures known in the art may be used to produce such antibodies and fragments. For preparation of monoclonal antibodies, any technique
25 which provides antibodies produced by continuous cell line cultures may be used, such as the hybridoma technique of Kohler and Milstein (*Nature*, 256:495-497 (1975)). Techniques for the production of single-chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to phosphoserine, phosphothreonine, or phosphotyrosine.

30 As may be seen from the above discussion, the assay format gives the user a great deal of latitude for tailoring reagents and reaction conditions to meet the requirements of a specific kinase or phosphatase.

Changes in the fluorescent properties of the reporter may be measured using any technique capable of detecting a change in the fluorescent properties of a molecule.
35 Such methods include, but are not limited to, fluorescence polarization, fluorescence quenching, fluorescence resonance energy transfer, and fluorescence correlati n

spectroscopy.

Fluorescence polarization: Fluorescence polarization (FP), in contrast to intensity measurements, can readily be used in the development of true, homogeneous, solution assays, or for real time, continuous recording assays, as the method can generate a direct quantitation of the ratio of bound/free ligand. This is extremely cost effective in terms of simplicity of operations and the types of screening laboratory ware required for the assay. Since polarization is independent of the fluorescence intensity, this technique can be used in the presence of colored compounds which may quench the emission.

When a fluorescent molecule is excited by polarized light, its emission is also polarized. The degree of polarization is dependent upon the viscosity of the solution, the rotational correlation time of the fluorophore (ϕ), and the temperature of the reaction mixture. Mathematically, the steady-state polarization (P) is expressed as

$$P = \frac{I_{\parallel} - I_{\perp}}{I_{\parallel} + I_{\perp}} \quad (2)$$

which is the difference between the intensities (I) of the parallel and perpendicular components of the polarized emission divided by their sum. More recently, the fluorescence anisotropy (r), which is dependent on the extent of rotational motion during the lifetime of the excited state, is being used because theoretical expressions are simpler when expressed in terms of this parameter rather than P. P is related to r by the expression

$$P = \frac{3r}{2 + r} \quad (3)$$

Since P, unlike r, is not completely linear with fraction bound, it is more beneficial to use r for the calculations, especially if the quantum yields of the free (q_F) and bound (q_B) species are not equal. The corrected fraction bound in this case is given by

$$FB = \frac{r - r_P}{(r_B - r)R + r - r_P} \quad (4)$$

where r_f and r_B are the anisotropies of the totally free and totally bound species respectively, r is the anisotropy of the experimentally observed bound species, and $R = q_B/q_F$. Titration of the fluorescent component with the nonfluorescent component yields data which is readily analyzable by various forms of the Langmuir binding isotherm or the stoichiometric binding equation if the binding is specific.

The requirements for a polarization assay are: (1) an assay component (the one with the lowest molecular weight) must be labelled with a long lifetime fluorescent probe and retain its ability to bind to the other component(s), (2) there must be a sufficient difference between the molecular weight of the labelled component and the nonlabeled one such that the probe senses a significant volume change upon binding, (3) the probe must have a relatively high quantum yield so that its fluorescence at low concentrations is significantly greater than background (4) the temperature and viscosity of the reaction mixture must be strictly controlled, and (5) the polarization should increase in a dose-dependent saturable manner. For use in plate-reader format, the fluorophore used should be a bright visible probe such as, e.g., fluorescein or rhodamine. Thus, where the detection method to be used is fluorescence polarization, the fluorescent label to be used may be any environment-sensitive probe whose fluorescence lifetime changes upon binding to a larger molecule.

The changes observed in the anisotropy free and bound ligands are a function of their individual rotational correlation times. Thus, in order to obtain a good dynamic range for the assay, the rotational correlation time of the labeled antigen should be shorter than the lifetime of the fluorescent tag. Since most visible probes have lifetimes ≤ 10 ns, the polarization assay is limited generally to the binding of small labelled ligands to large unlabelled targets. Thus it is generally not possible to study by polarization methods the interaction of two large macromolecules.

Fluorescence Quenching: HTS assays that utilize the intensity of the fluorophore as a readout usually require the separation of the free and bound species unless an environment-sensitive fluorophore is utilized. However, there are very few probes of this nature which both absorb and emit in the visible range, and as such, are not useful for HTS. In some cases, however, binding of the nonfluorescent component to the labelled ligand may result in a concentration-dependent quenching or enhancement of

the fluorescence which may occur due to the presence of quenching groups in the unlabelled component or some chemistry of the binding site which affects the fluorescence emission of the fluorophore. Thus the concentration-dependent decrease (or increase) in the fluorescence emission may be used to quantitate binding. The advantage of this readout is that a simple fluorescence plate reader may be used without polarizers. The usefulness of this assay may be limited where colored compounds which also quench the fluorophore are present in the assay mixture. Thus, where the detection method to be used is fluorescence quenching, the fluorescent label to be used may be any environment-sensitive probe whose fluorescence intensity changes upon binding to a larger molecule.

Fluorescence Resonance Energy Transfer: Fluorescence resonance energy transfer (FRET) is the transfer of the excited state energy from a donor (D) to an acceptor (A), and occurs only when the emission spectrum of the donor (D) fluorophore overlaps the absorption spectrum of the acceptor (A) fluorophore. Thus, by exciting at the absorption maximum of the donor and monitoring the emission at the long wavelength side of the acceptor fluorophore, it is possible to monitor only D and A molecules that are bound and reside within a certain distance, r . Thus one can monitor either the quenching of D or enhanced emission of A. The transfer rate, k_T in sec^{-1} is mathematically defined as

$$k_T = (r^{-6} J \kappa^2 n^{-4} \lambda_d) \times 8.71 \times 10^{23} \quad (5)$$

where r is the D-A distance in angstroms, J is the D-A overlap integral, κ^2 is the orientation factor, n is the refractive index of the media, and λ_d is the emissive rate of the donor. The overlap integral, J , is expressed on the wavelength scale by

$$J = \int_0^\infty F_d(\lambda) \epsilon_a(\lambda) \lambda^4 d\lambda \quad (6)$$

where its units are M^{-1}cm^3 , F_d is the corrected fluorescence intensity of the donor as a function of wavelength λ , and ϵ_a is the extinction coefficient of the acceptor in $\text{M}^{-1}\text{cm}^{-1}$.

Constant terms in equation 4 are generally combined to define the Forster critical distance, R_o , which is the distance in angstroms at which 50% transfer occurs. By substitution then, R_o can be defined in terms of the overlap integral, J , in angstroms, as

5

$$R_o = 9.79 \times 10^3 (\kappa^2 n^{-4} \phi_d J)^{1/6} \quad (7)$$

with ϕ_d being the quantum yield of the donor.

R_o and r are related to the transfer efficiency, E , by

10

$$E = \frac{R_o^6}{R_o^6 + r^6} \quad (8)$$

which determines the practical distance by which D and A can be separated to obtain a usable signal.

15 From these equations it is easy to see that, for high sensitivity, it is important to choose D-A pairs which have high quantum yields, high J values, and high R_o values. For example, R_o for the fluorescein/rhodamine pair is about 55Å. Large values of R_o are necessary to achieve a measurable signal when molecules containing D and A bind to each other. In practice it is common to use twice as
20 many acceptor as donor molecules in the reaction mixture if the emission of A is to be used as the readout. Thus, binding of, for example, macromolecule I labelled with D, to macromolecule II labelled with A, can be detected by the emission of A when excited at the absorption of D. Again, for a competitive process, the concentration increase in A fluorescence must occur in a hyperbolic, saturable
25 manner. The FRET assay is thus especially desirable for monitoring the binding of two large macromolecules where FP techniques are not particularly useful.

In view of the guidelines provided above and in the Examples, below, the skilled practitioner will be able to choose the detection technique best suited for the particular assay being performed.

30 Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of

illustration and are not intended as limiting.

EXAMPLES

5 *Example 1*

Materials and Methods

(a) *Reagents:* Antibodies to phosphoryltyrosine were obtained from ICN Biomedicals, Inc., Costa Mesa, CA, (monoclonal antibody clone PY20, Catalog Number 69-137), Life Technologies, Gaithersburg, MD, (monoclonal antibody clone 6G9, Catalog Number 13160-011), Molecular Biology Laboratories, Nagoya, Japan (monoclonal antibody clone 6D12; Catalog Number MH-11-3), Sigma, St. Louis MO, (monoclonal antibody clone PT-66, Catalog Number B-1531), Transduction Labs, Lexington, KY (monoclonal antibody clone PY20, Catalog Number P11120; altered Fab of PY20 produced in *E. coli*, Catalog Number E120H), Upstate Biologicals Inc., 15 Lake Placid NY, (monoclonal antibody clone 4G10, Catalog Number 05-321), and Zymed Laboratories, Inc., South San Francisco, CA, (monoclonal antibodies, clone PY-7E1, Catalog Number 13-5900; clone PY-IB2, Catalog Number 13-6300; clone PY20, Catalog Number 03-7700; clone Z027, Catalog Number 03-5800; and polyclonal antibodies Catalog Number 61-5800 and 61-8300).

20 Antibodies to phosphorylthreonine were obtained from Sigma (monoclonal antibody clone PTR-8, Catalog Number B-7661) and Zymed (monoclonal antibody, clone PT-5H5, Catalog Number 13-9200; polyclonal antibody, Catalog Number 61-8200). Antibodies to phosphorylserine were obtained from Sigma (monoclonal antibody clone PSR-45, Catalog Number B-7911) and Zymed (polyclonal antibody, 25 Catalog Number 61-8100). The structures of all chemicals used are shown in .

(b) *Synthesis of amino-fluoresceinated phosphorylamino acids:* The amino-fluoresceinated, phosphorylamino acids (i.e. - serine, threonine and tyrosine) were synthesized by reacting the phosphorylamino acids with fluorescein isothiocyanate (FITC) (Pierce Chemical Company) in 0.1 M NaHCO₃ of pH 9. The FITC used was a 30 mixture of the two isomers shown in Figures 1A and 1B. Purification was performed using a C4 Reverse Phase column (2.5 mm X 25 cm), a flow-rate of 0.5 ml/min, and a 15 minute gradient from 0.01 N HCl to 100% CH₃CN. The fluoresceinated phosphorylamino acids were identified by mass spectral analysis. For these compounds, a molar extinction coefficient of 77,000 was assumed in pH 8 35 buffer with a 1 cm pathlength.

(c) *Fluorescence Measurements:* Ratiometric fluorescence measurements were

made using an ISS K2 spectrofluorometer equipped with Glans Thompson quartz polarizers. The temperature of the reaction cuvette was maintained at 25.0°C by a Lauda RM6 circulating bath. The fluorescence of the labelled phosphorylamino acids was measured with excitation at 485 nm and emission was observed through an

5 Omega 530 ± 15 nm bandpass filter.

Antibody binding to the fluoresceinated phosphorylamino acids was monitored by adding successive small volumes of the antibody stock solution to a cuvette containing a fixed amount of fluoresceinated phosphorylamino acid in two milliliters of pH 7.4 buffer (25 mM Tris and 25 mM NaCl adjusted to pH 7.4 with

10 HCl). The fluorescence polarization, anisotropy, and the intensity — the sum of vertically and horizontally polarized emissions — were recorded one minute after each addition. The dilution produced by the addition of the antibody necessitated a slight correction for both the concentrations and the intensity readings but not for the polarization since the latter is independent of the fluorophore concentration.

15 Displacement of the fluorophore from the antibody by competing nonfluorescent ligands was measured by using fixed amounts of antibody and fluorophore in two milliliters of pH 7.4 buffer and adding successive small volumes of the competitor stock solution. Again, the displacement was monitored one minute after each addition by the changes in fluorescence intensity, polarization, and anisotropy

20 *Data Analysis:* Data analysis was performed as follows. It is assumed that the binding of the fluorophore ligand, L, and the competing inhibitor, I, to the antibody, A, are rapid, simple thermodynamic equilibria, according to the scheme:



25



with $K_1 = L/A/C$ and $K_2 = I/A/X$.

Case I: Displacement with $A_0 \gg L_0$: For the displacement experiments —

30 where both L_0 and A_0 are constant and I_0 increases — the analysis of the data is

greatly simplified when it is possible to use experimental conditions where $I_o \gg A_o$ and $A_o \gg L_o$. Under these conditions, $L = L_o - C$; $I = I_o$; $X = A \cdot I_o / K_i$; $A_o = A + X = A(1 + I_o / K_i)$. Therefore $A = A_o / (1 + I_o / K_i)$ and

$$K_i = \frac{(L_o - C) A_o}{C \cdot \left(1 + \frac{I_o}{K_i}\right)} \quad (3)$$

5

Upon solving (3) for C we obtain:

$$C = \frac{C_u}{1 + \frac{I_o}{K_i^{app}}} \quad (4)$$

10

in which we define the concentration of the fluorophore/antibody complex in the absence of inhibitor as $C_u = L_o / (1 + K_i / A_o)$ and the apparent inhibition constant as $K_i^{app} = K_i \cdot (1 + A_o / K_i)$. In these experiments one can directly determine only K_i^{app} . The value of K_i can be calculated only if L_o and K_i are known.

15

Case IIa: Binding: The analysis of the direct binding of a constant concentration of fluorophore, L_o , to a variable concentration of antibody, A_o , of comparable magnitude requires solving a quadratic equation. By substituting L and A from the stoichiometric equations $L_o = L + C$ and $A_o = A + C$ into the definition of K_i we obtain:

20

$$K_i = \frac{(L_o - C) \cdot (A_o - C)}{C} \quad (5)$$

which yields the quadratic equation:

$$C^2 - C \cdot (A_o + L_o + K_I) + L_o A_o = 0 \quad (6)$$

The solution of (6) is

$$C = \frac{1}{2} \cdot (A_o + L_o + K_I - \sqrt{(A_o + L_o + K_I)^2 - 4 \cdot A_o \cdot L_o}) \quad (7)$$

5

Case IIb: Displacement: The usual conditions for the displacement experiments are L_o and A_o constant but of comparable magnitude with increasing I_o and $I_o \gg A_o$. Under these conditions, $L = L_o - C$; $I = I_o$; and $X = A \cdot I_o / K_I$. Therefore A_o

10 $= A + C + X = C + A \cdot (1 + I_o / K_I)$ so that

$$A = \frac{A_o - C}{1 + \frac{I_o}{K_I}} \quad (8)$$

and

15

$$K_I C = L \cdot A = \frac{(L_o - C) \cdot (A_o - C)}{1 + \frac{I_o}{K_I}} \quad (9)$$

Solving (9) for C yields

$$C = \frac{1}{2} \cdot \left(A_o + L_o + K_I \cdot \left(1 + \frac{I_o}{K_I} \right) - \sqrt{\left[A_o + L_o + K_I \cdot \left(1 + \frac{I_o}{K_I} \right) \right]^2 - 4 \cdot A_o \cdot L_o} \right) \quad (10)$$

20

Fluorescence Intensity: The fluorescence intensity, F , of a mixture of L and C is

$$F = Q_L \cdot L + Q_C \cdot C = Q_L \cdot L_0 + (Q_L - Q_C) \cdot C \quad (11)$$

5

where Q_L is the molar emissivity of the free fluorophore and Q_C that of the bound one. Substituting into this equation C from Eq 7 yields the quadratic used to analyze the fluorescence intensity binding data obtained under Case IIa.

Similarly, substituting the expression for C from Eq 4 or from Eq 10 yields the expression for analyzing the fluorescence displacement data under Cases I or IIb, respectively.

Fluorescence Polarization and Anisotropy: The analysis of the changes in fluorescence intensity upon binding to the antibody are relatively simple because the changes are a linear function of the composition of the $L+C$ mixture. As pointed out previously (7), this is not the case in general when changes in fluorescence polarization or anisotropy are measured. Indeed, if the exciting electric vector is vertical, then the polarization of a solution is a function of the emitted intensities horizontally polarized, h , and vertically polarized, v , according to the definition:

$$p = \frac{v - h}{v + h} \quad (12)$$

20

The total fluorescence intensity is $F = h + v = Q \cdot M$, where Q is the molar emissivity of the fluorophore at concentration M . Solving Eq 12 for v yields $v = h \cdot (1+p)/(1-p)$.

Thus:

25

$$Q \cdot M = h \cdot \left(1 + \frac{1+p}{1-p} \right) \quad (13)$$

and it follows that

$$h = \frac{Q \cdot M (1 - p)}{2} \quad (14)$$

with

$$v = \frac{Q \cdot M (1 + p)}{2} \quad (15)$$

5

For a mixture of L and C, the polarizations calculated from the sum of the vertically and horizontally polarized emission intensities are:

$$p = \frac{v_L + v_C - h_L - h_C}{v_L + v_C + h_L + h_C} \quad (16)$$

10

Substituting $h_L = Q_L L(1-p_L)/2$, $h_C = Q_C C(1-p_C)/2$, $v_L = Q_L L(1+p_L)/2$, and $v_C = Q_C C(1+p_C)/2$, where p_C and p_L are the polarization values for the bound ligand and the free ligand, respectively, and then solving for C/L yields, as reported earlier (7):

15

$$\frac{C}{L} = \frac{C}{L_0 - C} = \frac{Q_L}{Q_C} \frac{p - p_L}{p_C - p} \quad (17)$$

Similarly, from the definition of the fluorescence anisotropy, a,:

$$a = \frac{v - h}{v + 2h} \quad (18)$$

20

we derive:

$$\frac{C}{L} = \frac{C}{L_o - C} = \frac{Q_L}{Q_C} \frac{2 + a_C}{2 + a_L} \frac{a - a_L}{a_C - a} \quad (19)$$

where a_C and a_L are the fluorescence anisotropies for the bound ligand and the free ligand, respectively.

Eq 17 can be solved for p as a function of C :

5

$$p = \frac{p_L L_o + C \left[\frac{Q_C}{Q_L} p_C - p_L \right]}{L_o + C \left[\frac{Q_C}{Q_L} - 1 \right]} \quad (20)$$

Of course, when the molar emissivity does not change upon binding, i.e. when $Q_C/Q_L = 1$, then p becomes a linear function of C , in full analogy with the
10 fluorescence intensity as expressed in Eq 11.

When $Q_C/Q_L \neq 1$ then substituting into Eq 20 the value of C from the appropriate equation (Eq 4, Eq 7, or Eq 10) allows one to determine K_i and K_i^{app} or K_i by nonlinear least squares analysis of the polarization data. Depending on the experimental conditions used, such an analysis may also yield the best-fit values for
15 p_L , p_C , Q_L , and Q_C .

Results

The majority of the anti-phosphorylamino acid antibodies available from suppliers were evaluated, looking for those that produced changes in either the
20 intrinsic polarization or quenching of the emission intensity of the fluoresceinated phosphorylamino acids. From the results of this survey, three classes of antibodies were identified for phosphoryltyrosine, based on the types of fluorescent signal produced by the antibody: (I) those giving a large polarization change without significant effect on the fluorescence emission of the fluorescent phosphorylamino
25 acid, (II) those producing polarization changes and quenching of the emission, and (III) those yielding little or no change in either parameter, or produced noisy, nonreproducible data.

Due to sensitivity considerations, titrations of the fluorescent

phosphorylamino acids with the antibodies were performed at comparable concentrations of A_0 and L_0 . Under these conditions, one must use the quadratic form of the binding equation (Equation 7) in order to determine with precision the K_d of the ligand for the antibody and the stoichiometry of the reaction. The concentration of L_0 was calculated from the extinction coefficient of the fluorophore. However, the concentration of the stock solution of the antibody is given in units of protein concentration/ml, not active antibody which has two binding sites/molecule.

The antibody concentrations were first calculated using the protein concentrations supplied by the manufacturer, assuming a molecular weight of 160,000. [1]. Calculations done in this manner do not take into consideration the bidentate nature of the antibody. Thus, the site-concentration or normality of the antibody should be twice the concentration calculated as described above. Calculation of antibody concentrations based on protein quantitation would exceed the actual antibody concentration, due to the presence of extraneous contaminating proteins and/or inactive antibody. For these reasons, in the titration curves, antibody concentrations are given in terms of the volume of antibody added, and the actual K_d s of the fluorescent phosphorylamino acids are calculated from the fitted stoichiometry, which has concentration units of molar sites (i.e., normality). Comparison of the determined stoichiometry with concentrations calculated from protein concentrations can then be used to assess the purity and activity of antibodies from different manufacturers.

Examples of experimental results using Type I antibodies (as described above) are shown in Figures 2A-2C. Addition of any of the Sigma anti-phosphorylamino acid antibodies to the corresponding fluoresceinated phosphorylamino acid produced a steep, concentration-dependent increase in the polarization of the fluorescence which was saturable at higher antibody concentrations. The maximal increase in polarization at saturating antibody concentrations was approximately 9-fold for antibody binding to labelled (i.e. - fluoresceinated) phosphorylamino acids with an excellent signal-to-noise ratio. The Sigma antibodies are thus Type I antibodies (FP change only). The data sets were analyzed using Equation 7 in conjunction with a nonlinear least squares fitting program and are consistent with this model as evidenced by the agreement between the experimental data points and the theoretical curves. The dissociation constant for all the Sigma antibodies, calculated from the fits to the experimental data, are given in Table 1 in units of antibody normality. All three fluoresceinated phosphorylamino acids had a high picomolar to low nanomolar affinities for their

corresponding antibodies.

The antibodies were then individually evaluated for their cross-reactivity with the opposite fluoresceinated phosphorylamino acids, and the non-phosphorylated amino acids themselves by competitive displacement immunoassay.

- 5 None of the antibodies were found to cross react with the other phosphorylamino or amino acids at levels 10-20 fold above those used in the titration experiments. The data for the anti-phosphoserine and anti-phosphothreonine antibodies are discussed more extensively below.

- 10 Three anti-phosphotyrosine antibodies which gave both significant increases in the fluorescence polarization and decreases in the emission intensity when bound to P-Tyr-F were also identified (Type II antibodies). Two antibodies, from ICN and MBL, produced the largest decreases in the fluorescence of P-Tyr-F upon binding (Table 1; as Q/Q_0 increases so does the dynamic range).

- 15 Analysis of the data in Figures 3A and 3B and 4A-4C using Equations 7 and 20 resulted in fits that were consistent with the experimental points, and there was good agreement between the K_d s (low nM) calculated from both the polarization and the intensity readings. Thus, kinase and phosphatase assays utilizing the ICN or MBL antibodies may be performed using either the fluorescence polarization increase or emission quenching as a measure of antibody binding to labeled
20 phosphorylamino acids in the presence of unlabeled phosphorylated reaction products. The cause of the emission quenching by the ICN and MBL antibodies may involve hydrogen bonding of the fluorescent ligand, the presence of tryptophan residues in or near the binding site, or electrostatic interactions.

- Other antibodies were found to produce small, nonrobust fluorescence
25 changes upon binding labelled ligand (Table 1), and were considered not to be useful for the kinase assays. Still other antibodies tested produced insignificant or inconsistent changes in either the emission intensity or polarization, and were deemed not usable for either kinase or phosphatase activity detection (Type III antibodies). Interestingly, the superior antibodies (those that gave the most robust
30 signals with the least scatter) had normalities calculated from the fits to the quadratic equation that were in good agreement with concentrations supplied by the manufacturer. In contrast, for the less robust antibodies, there was a great disparity between the stoichiometry and protein concentration calculations, indicating the presence of either contaminating proteins and/or inactive antibody.

- 35 Antibodies for phosphoserine and phosphothreonine came from two sources, Sigma (St. Louis, MO) and Zymed (CA). Only the Sigma antibodies were found to


produce a signal of significant magnitude that would allow, for kinase and phosphatase assays, accurate quantitation of unlabelled phosphoserine and phosphothreonine (Table 2). The results from experiments with the Sigma antibodies were calculated by nonlinear least squares analysis using equation 7, are
5 shown in Figures 2A-2C. None of the Sigma antibodies produced quenching of the emission intensities, but both P-Ser-F and P-Thr-F had high affinity for their respective antibodies as measured by FP. Although binding of the fluorescent ligands to the Zymed antibodies gave both polarization and intensity changes, they were accompanied by significant scatter and a less than desirable dynamic range to
10 be useful for monitoring either kinase or phosphatase activity.

In order to determine if kinase reactions can be followed by competition of the phosphorylated substrate with labelled phosphorylamino acid, the specificity of the binding by the labeled phosphorylamino acids was demonstrated by
co-competition with their unlabeled counterparts for the antibody-bound fluorescent
15 complex. The competition between the labeled and unlabeled phosphorylamino acids for the antibodies was first determined. The amounts of antibody and fluorescent phosphorylamino acid were held constant (close to the K_d) in these experiments, and the decrease in polarization or increase in emission resulting from the addition of the unlabelled phosphorylamino acid was monitored as a function of its
20 concentration. The experimental data were analyzed by fixing the concentrations of L , K_d , and Q_d using Equation 10 in conjunction with Equation 20 (when polarization with a quenching antibody was measured), or Equation 10 only when the increased polarization was not accompanied by quenching. The results of these experiments are shown in Figures 4A-4C (Sigma antibodies) and 5A and 5B (MBL antibody).
25 Fitting in the manner described above yielded excellent fits to the experimental data. The values for the corrected dissociation constants of the phosphorylamino acids are given in Table 3. The K_d s for phosphoryltyrosine, phosphorylserine, and phosphorylthreonine were considerably higher than those for the fluoresceinated ligands. Good agreement between K_d s calculated from the polarization and
30 quenching data were determined for the MBL antibody. These experiments show that kinase and phosphatase activities can be measured with precision using either FP or quenching data.

In order to calculate, by competition with labeled ligand, the amount of phosphorylated or dephosphorylated substrates produced during the course of the
35 kinas or phosphatase reactions, it is imperative to know the dissociation constant of the phosphorylated substrate for the particular antibody. The affinities of tw


phosphorylated peptides which are substrates for the JAK-2 tyrosine kinase were measured by displacement of fluoresceinated phosphoryltyrosine from both the corresponding Sigma and MBL antibodies. The structure of the first JAK-2 substrate, JAK-2(a), is:

5



10 The formula of the second JAK-2 substrate, JAK-2(b) is:

15




Both phosphorylated peptides displaced P-Tyr-F from the antibodies in a dose-dependent manner, whether measured by FP only (Sigma) or simultaneously using quenching or fluorescence polarization (MBL). The data were analyzed by nonlinear least squares analysis using equation 7, and the results are shown in Figures 6 and 7A-7D. The K_s calculated from the fits are given in Table 3. The K_s of the two phosphorylated JAK-2 kinase substrates for Sigma antibody were one and one-half orders of magnitude higher than those of the labelled ligands. Similar results were obtained for displacement from the MBL antibody as shown in Figures 7A-7D, and the calculated K_s for this antibody from both FP and quenching measurements were in good agreement. Thus, using the fluoresceinated phosphorylamino acids in a competition reaction to measure kinase and phosphatase activities requires that the concentrations of added fluoresceinated ligand should be comparable to K_d and the amount of A up to $10 \times L$. Under these degenerate conditions, the data can then be analyzed by Equation 4.

Discussion

35 Two superior anti-phosphotyrosine antibodies were identified that produced changes in the fluorescence polarization of a bound ligand or quenching of its

emission intensity, and would enable quantitation of the activities of any phosphatase or kinase. Kinase phosphorylation of a substrate peptide or protein was monitored by specific displacement of a fluorescent phosphorylamino acid or a fluorescent phosphorylated peptide from an antibody by the reaction product from the kinase assay (see reaction scheme below). A proposed assay format for the fluorescent kinase assay is given below.



- 10 In this scheme, the kinase reaction proceeds for a period of time, is stopped by the addition of EDTA, and both antibody and fluoresceinated phosphorylamino acid added. Alternatively the antibody and labeled ligand could be present at time = 0. The phosphorylated substrate would compete with the fluoresceinated phosphorylamino acid for the antibody, giving intermediate values of polarization and/or quenching. Phosphatase activity can also be monitored using this immunoassay. Several fluorescent substrates have previously been used to kinetically follow phosphatase activity. These include phosphotyrosine (Z.Y. Zhang *et al.*, *Anal. Biochem* 211:7-15 (1993); B. Galvan *et al.*, *Clin. Biochem.* 29: 125-131 (1996)), 2-methoxybenzoyl phosphate (P. Paoli *et al.*, *Experientia* 51:57-62 (1995)), europium-labeled antibody (D. Worm, *Diabetologia* 39: 142-148 (1996)); terbium chelates (T.K. Christopoulos, *Anal. Chem.* 64: 342-346 (1992)), and fluorescein diphosphate (E. Tolsa *et al.*, *J. Immunol. Methods* 192: 165-172 (1996)), with fluorescein diphosphate being the most sensitive and amenable to HTS.
- 20 The evaluated antibodies demonstrated high affinities for the fluoresceinated phosphoryl amino acids. Therefore, a significant concentration of phosphorylated peptide would be required to displace the labeled ligand. This is detrimental for

kinase assays. However, this problem can be obviated by selecting three generic peptides, one for each of the three amino acids, that is phosphorylated and fluorescently labeled. Peptides having reduced affinity for the antibodies will allow detection of lower concentrations of phosphorylated product from the kinase reactions. With proper selection of the peptide and its fluorescent labeling, the assay format can be FP, FRET, FQ, or FCS.

Two anti-phosphoryltyrosine antibodies (Sigma, St. Louis, MO, and MBL International Corp., Watertown, MA) were identified that produced robust fluorescence signals upon binding a labeled phosphoryltyrosine or phosphoryltyrosine peptide. The MBL antibody not only yielded a large polarization change, but also significantly quenched (*i.e.*, reduced by 50%) the probe fluorescence. By using an antibody-reporter pair which, upon binding to each other, exhibit a change in both FP and FQ, the kinase assay described above can be performed in any fluorescent plate reader, not just those with polarization capabilities. The use of polarization as a readout is desirable since this parameter is independent of the fluorescence emission intensity (Leavine, L.M., *et al.*, *Anal. Biochem.* 247:83-88 (1997)) and is thus not subject to the optical artifacts imparted by colored compounds which may quench the fluorescence emission of the label. Several other anti-phosphoryltyrosine antibodies from other suppliers were also tested (results not shown but the suppliers are listed in Materials and Methods) and found to produce low or inconsistent fluorescence changes upon binding the fluorescent ligand.

Four antibodies against phosphoserine and phosphothreonine (obtained from Sigma and Zymed) were tested. The Sigma antibodies against phosphoserine and phosphothreonine produced fluorescence polarization changes of sufficiently large dynamic range that the antibodies can easily be used to measure the activities of specific kinases.

All three fluorescein labelled phosphorylamino acids had higher affinities for their specific antibodies than their unlabeled counterparts, presumably due to their relatively small size and the increased hydrophobicity contributed by the fluorescein moiety. Due to the high affinity of the antibodies for the labeled phosphorylamino acids, a significant amount of product must be generated by kinase activity before displacement can occur. However, any labeled phosphorylated peptide or small protein, which would have lower affinity for the antibody could instead be used as the competitive ligand, including fluorescein-modified phosphorylated JAK-2 substrate peptides. Thus, the assay format gives the user a great deal of latitude for tailoring reagents and reaction conditions to meet the requirements of a specific

kinase or phosphatase.

Example 2: Identification of peptides having a decreased affinity for Antibody
Materials and Methods

- 5 *Fluorescent kinase assay:* As is discussed above, the displacement of reporter molecule by phosphorylated substrate in the competitive immunoassays of the invention will occur at lower concentrations of phosphorylated substrate, and thus most amenable to HTS, where the K_d of the antibody for the phosphorylated substrate is less than or equal to the K_d of the antibody for the reporter molecule.
- 10 This may be accomplished by obtaining a small phosphorylated labeled peptide which has a lower binding affinity toward its antibody than the corresponding fluorescent phosphorylated amino acid (between about 0.5 nM and 1 nM for fluoresceinated phosphotyrosine or phosphoserine). Appropriate peptides may have between about 3 and 50 amino acid residues, preferably between about 3 and 25
- 15 residues, more preferably between about 3 and 15 residues, and most preferably between about 4 and 10 amino acid residues. Such peptides may be chemically synthesized, may be the result of enzymatic or chemical cleavage of a larger peptide or protein, or may be produced recombinantly. The peptides are then labeled and purified, and their affinities toward their corresponding antibodies are measured.
- 20 Adjustments to the length and sequence of the peptides can be made to the peptides if this is deemed necessary as a result of their K_d for antibody. Pentapeptides have been prepared which contain phosphorylated amino acids with a free cysteine available for labeling with a fluorophore. Peptides having the desired affinity (between about 50 nM and 100 nM) are then tested in the assay system for detecting
- 25 the phosphorylation of substrate by kinases. As the assay is only measuring product, it will be possible to identify the actual amino acid of a substrate that is being phosphorylated by a kinase using cocompetition experiments. For example, a peptide substrate that has been phosphorylated by a kinase can be used to displace reporter peptides containing individual phosphorylated amino acids (i.e., either Ser,
- 30 Thr or Tyr) from their respective antibodies.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

Numerous modifications and variations of the present invention are possible

35 in light of the above teachings and, therefore, are within the scope of the invention. The entire disclosure of all publications cited herein are hereby incorporated

by reference.

**Table 1. Affinities of Anti-phosphoryltyrosine Antibodies for
Fluoresceinated Phosphoryltyrosine**

Antibody	Fluorescence Polarization		Fluorescence Quenching	
	K_d , nM sites (p) ¹	Δp ¹	K_d , nM Sites (Φ) ²	Q/Q_0 ³
Sigma	1.9 ± 0.3	x9	-----	-----
Gibco BRL	9.8 ± 7.3	x6	-----	-----
ICN	3.8 ± 1.0	x6	5.1 ± 1.6	1.55
UBI	bad data	x5	81.8 ± 52.1	1.55
MBL	4.7 ± 1.1	x9	3.9 ± 0.3	2.05
Zymed rabbit polyclonal	6.6 ± 1.3	x8	1.7 ± 0.7	1.28
Zymed PY Plus	poor signal	x2	4.2 ± 2.7	1.1

¹ Direct polarization measurements

² Φ is the decreased Quantum Yield used to calculate the K_d

³ Ratio of the Quantum Yields of free and bound probe. Used to correct polarization measurements as described in Data Analysis.

5 **Table 2. Affinities of Anti-phosphorylserine and Anti-phosphorylthreonine
Antibodies for Fluoresceinated Phosphorylamino Acids**

Antibody	Fluorescence Polarization	
	K_d , nM Sites (p)	Δp
Sigma Anti P-Ser	0.30 ± 0.03	x9
Zymed Anti P-Ser	$\approx 8.13^*$	x2.5
Sigma Anti P-Thr	1.055 ± 0.9	x5.1
Zymed Anti P-thr	58.1 ± 10.1	x5

* Signal too noisy for accurate quantitation

Table 3. Affinities of Unlabelled Phosphorylamino Acids and Phosphorylated JAK-2 Substrates for their Respective Antibodies¹

5	Phosphoryl Ligand	Sigma Monoclonal		MBL Monoclonal	
		K_d , nM (p)	K_d , nM ϕ	K_d , nM (p)	K_d , nM ϕ
	Phosphoryltyrosine	11.9 ± 2.5	----	5.5 ± 1.0	5.4 ± 0.5
	Phosphorylserine	16.8 ± 3.1	----	----	----
10	Phosphorylthreonine	44.1 ± 6.0	----	----	----
	JAK-2(a)	15.4 ± 1.6	----	8.1 ± 1.8	11.6 ± 4.1
	JAK-2(b)	49.5 ± 3.6	----	12.1 ± 1.1	8.1 ± 0.9

15 ¹ K_d s are calculated for nM sites (i.e. - normality)

What is claimed is:

1. A method of determining the phosphorylating activity of an enzyme comprising the steps of:
 - 5 (a) combining said enzyme with
 - (i) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein said amino acid is selected from the group consisting of serine, threonine and tyrosine;
 - (ii) a substrate molecule comprising the same amino acid that is
10 phosphorylated in said reporter, wherein said substrate molecule is capable of being phosphorylated at said amino acid by said enzyme to yield a product;
 - (iii) an antibody which selectively binds to a molecule comprising said phosphorylated amino acid; and
 - (iv) a phosphate source;
 - 15 (b) measuring the FP, FQ, or FCS of said reporter following the combination of step (a); and
 - (c) using the FP, FQ, or FCS measurement of step (b) to determine the activity of said enzyme.
- 20 2. The method of claim 1, wherein said substrate is combined with said enzyme before the addition of said reporter and said antibody.
3. The method of claim 1, wherein said substrate, said reporter, and said antibody are combined with said enzyme simultaneously.
- 25 4. The method of claim 1, wherein the K_d of the product for the antibody is less than or equal to the K_d of the antibody for the reporter molecule.
5. The method of claim 4, wherein the K_d of the product for the antibody is less
30 than the K_d of the antibody for the reporter molecule.
6. The method of claim 1 wherein the fluorescence polarization of said reporter is measured in step (b).
- 35 7. The method of claim 1 wherein the fluorescence quenching of said reporter is measured in step (b).

8. The method of claim 1 wherein the FCS of said reporter is measured in step (b).
9. The method of claim 1, wherein the substrate molecule is phosphorylated at a serine residue.
- 5 10. The method of claim 1, wherein the substrate molecule is phosphorylated at a threonine residue.
11. The method of claim 1, wherein the substrate molecule is phosphorylated at a
10 tyrosine residue.
12. The method of claim 11, wherein said reporter molecule is a peptide having between about 4 and 15 amino acid residues.
- 15 13. A method of screening for an agent capable of increasing or decreasing the phosphorylating activity of an enzyme comprising the steps of:
- (a) performing the method of claim 1 in the presence and in the absence of said agent;
- (b) comparing the activity of said enzyme in the presence of said agent with
20 the activity of said enzyme in the absence of said agent to determine whether the phosphorylating activity of said enzyme in the presence of said agent is increased or decreased.
14. A method for determining the dephosphorylating activity of an enzyme
25 comprising the steps of:
- (a) combining said enzyme with
- (i) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein said amino acid is selected from the group consisting of serine, threonine and tyrosine;
- 30 (ii) a substrate molecule comprising the same phosphorylated amino acid as said reporter, wherein said substrate molecule is capable of being dephosphorylated at said amino acid by said enzyme to yield a product;
- (iii) an antibody which selectively binds to a molecule comprising said phosphorylated amino acid; and
- 35 (iv) a phosphate acceptor;
- (b) measuring the FP, FQ, or FCS of said reporter following the combination

of step (a); and

(c) using the FP, FQ, or FCS measurement of step (b) to determine the activity of said enzyme.

5 15. The method of claim 14, wherein said substrate is combined with said enzyme before the addition of said reporter and said antibody.

16. The method of claim 14, wherein said substrate, said reporter, and said antibody are combined with said enzyme simultaneously.

10

17. The method of claim 14, wherein the K_d of the product for the antibody is less than or equal to the K_d of the antibody for the reporter molecule.

18. The method of claim 17, wherein the K_d of the product for the antibody is less
15 than the K_d of the antibody for the reporter molecule.

19. The method of claim 14 wherein the fluorescence polarization of said reporter is measured in step (b).

20 20. The method of claim 14 wherein the fluorescence quenching of said reporter is measured in step (b).

21. The method of claim 14 wherein the FCS of said reporter is measured in step (b).

25

22. The method of claim 14, wherein the substrate molecule is phosphorylated at a serine residue.

23. The method of claim 14, wherein the substrate molecule is phosphorylated at a
30 threonine residue.

24. The method of claim 14, wherein the substrate molecule is phosphorylated at a tyrosine residue.

35 25. A method of screening for an agent capable of modulating the dephosphorylating activity of an enzyme comprising the steps of:

- (a) performing the method of claim 14 in the presence and in the absence of said agent;
- (b) comparing the activity of said enzyme in the presence of said agent with the activity of said enzyme in the absence of said agent to determine whether the dephosphorylating activity of said enzyme in the presence of said agent is increased or decreased.

26. A method for determining the phosphorylation of a substrate molecule by an enzyme at an amino acid selected from the group consisting of serine, threonine and tyrosine, said method comprising the steps of:

- (a) combining said substrate molecule with
- (i) said enzyme
 - (ii) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein said amino acid is selected from the group consisting of serine, threonine and tyrosine;
 - (iii) an antibody which selectively binds to a molecule comprising said phosphorylated amino acid; and
 - (iv) a phosphate source;
- (b) measuring the FP, FQ, or FCS of said reporter following the combination of step (a); and
- (c) using the FP, FQ, or FCS measurement of step (b) to determine whether said substrate molecule has been phosphorylated.

27. A method for determining the dephosphorylation of a substrate molecule by an enzyme, wherein said substrate molecule comprises a phosphorylated amino acid, and wherein said amino acid is selected from the group consisting of serine, threonine and tyrosine, comprising the steps of:

- (a) combining said substrate molecule with
- (i) said enzyme
 - (ii) a reporter molecule comprising a fluorescent label and a phosphorylated amino acid, wherein said reporter molecule comprises the same phosphorylated amino acid as said substrate molecule;
 - (iii) an antibody which selectively binds to a molecule comprising said phosphorylated amino acid; and
 - (iv) a phosphate acceptor;
- (b) measuring the FP, FQ, or FCS of said reporter following the combination

of step (a); and

(c) using the FP, FQ, or FCS measurement of step (b) to determine whether said substrate molecule has been dephosphorylated.

5 28. A method of determining the phosphorylating activity of an enzyme comprising the steps of:

(a) combining the enzyme with:

(i) a substrate molecule comprising an amino acid selected from the group consisting of Ser, Thr, and Tyr, wherein said substrate molecule is capable of
10 being phosphorylated at said amino acid by said enzyme to yield a product, and wherein said substrate molecule is labeled with an acceptor fluorophore;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with a donor fluorophore which corresponds to the acceptor fluorophore labeling said substrate; and

15 (iii) a high-energy phosphate source;

(b) measuring the FRET of the combination of step (a); and

(c) using the FRET measurement of step (b) to determine the activity of the enzyme.

20 29. A method of determining the phosphorylating activity of an enzyme comprising the steps of:

(a) combining the enzyme with:

(i) a substrate molecule comprising an amino acid selected from the group consisting of Ser, Thr, and Tyr, wherein said substrate molecule is capable of
25 being phosphorylated at said amino acid by said enzyme to yield a product, and wherein said substrate molecule is labeled with a donor fluorophore;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore which corresponds to the donor fluorophore labeling said substrate; and

30 (iii) a high-energy phosphate source;

(b) measuring the FRET of the combination of step (a); and

(c) using the FRET measurement of step (b) to determine the activity of the enzyme.

35 30. A method of determining the dephosphorylating activity of an enzyme comprising the steps of:

- (a) combining the enzyme with:
- (i) a substrate molecule comprising a phosphorylated amino acid selected from the group consisting of phosphoserine, phosphothreonine and phosphotyrosine, wherein said substrate molecule is labeled with an acceptor fluorophore;
 - 5 (ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with a donor fluorophore which corresponds to the acceptor fluorophore labeling said substrate; and
 - (iii) a high-energy phosphate source;
- (b) measuring the FRET of the combination of step (a); and
- 10 (c) using the FRET measurement of step (b) to determine the activity of the enzyme.

31. A method of determining the dephosphorylating activity of an enzyme comprising the steps of:

- 15 (a) combining the enzyme with:
- (i) a substrate molecule comprising a phosphorylated amino acid selected from the group consisting of phosphoserine, phosphothreonine and phosphotyrosine, wherein said substrate molecule is labeled with a donor fluorophore;
 - (ii) an antibody which selectively binds to a molecule comprising the
 - 20 phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore which corresponds to the donor fluorophore labeling said substrate; and
 - (iii) a high-energy phosphate source;
- (b) measuring the FRET of the combination of step (a); and
- (c) using the FRET measurement of step (b) to determine the activity of the
- 25 enzyme.

32. A method of screening for an agent capable of modulating the phosphorylating or dephosphorylating activity of an enzyme comprising the steps of:

- (a) performing the method of claim 28, 29, 30, or 31 in the presence and in
- 30 the absence of said agent;
- (b) comparing the activity of said enzyme in the presence of said agent with the activity of said enzyme in the absence of said agent to determine whether the phosphorylating activity or the dephosphorylating activity of said enzyme in the presence of said agent is increased or decreased.

35

33. A method for determining the phosphorylation of a substrate molecule by an

enzyme at an amino acid selected from the group consisting of serine, threonine and tyrosine, wherein said substrate molecule is labeled with an acceptor fluorophore, comprising the steps of:

- (a) combining the substrate molecule with
 - 5 (i) the enzyme
 - (ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with a donor fluorophore which corresponds to the acceptor fluorophore labeling said substrate; and
 - (iii) a high-energy phosphate source;
- 10 (b) measuring the FRET of the reporter following the combination of step (a);
- and
- (c) using the FRET measurement of step (b) to determine whether the substrate molecule has been phosphorylated.

- 15 33. A method for determining the phosphorylation of a substrate molecule by an enzyme at an amino acid selected from the group consisting of serine, threonine and tyrosine, wherein said substrate molecule is labeled with a donor fluorophore, comprising the steps of:

- (a) combining the substrate molecule with:
 - 20 (i) the enzyme
 - (ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore which corresponds to the donor fluorophore labeling said substrate; and
 - (iii) a high-energy phosphate source;
- 25 (b) measuring the FRET of the reporter following the combination of step (a);
- and
- (c) using the FRET measurement of step (b) to determine whether the substrate molecule has been phosphorylated.

- 30 34. A method for determining the dephosphorylation of a substrate molecule by an enzyme, wherein the substrate molecule comprises a phosphorylated amino acid selected from the group consisting of phosphoserine, phosphothreonine and phosphotyrosine, and wherein said substrate molecule is labeled with an acceptor fluorophore comprising the steps of:

- 35 (a) combining the substrate molecule with:
 - (i) the enzyme;

(ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with a donor fluorophore which corresponds to the acceptor fluorophore labeling said substrate;

(b) measuring the FRET of the reporter following the combination of step (a);

5 and

(c) using the FRET measurement of step (b) to determine whether the substrate molecule has been dephosphorylated.

35. A method for determining the dephosphorylation of a substrate molecule by an
10 enzyme, wherein the substrate molecule comprises a phosphorylated amino acid selected from the group consisting of serine, threonine and tyrosine, and wherein said substrate molecule is labeled with a donor fluorophore comprising the steps of:

(a) combining the substrate molecule with:

(i) the enzyme;


15 (ii) an antibody which selectively binds to a molecule comprising the phosphorylated amino acid, said antibody being labeled with an acceptor fluorophore which corresponds to the donor fluorophore labeling said substrate;

(b) measuring the FRET of the reporter following the combination of step (a);


and

20 (c) using the FRET measurement of step (b) to determine whether the substrate molecule has been dephosphorylated.


FIGURES 1A - 1G




5-Fluorescein Isothiocyanate (Isomer 1)




6-Fluorescein Isothiocyanate (Isomer II)




5-Fluorescein Phosphoryl Amino Acid (Isomer I Derivative)




6-Fluorescein Phosphoryl Amino Acid (Isomer II)



Phosphoryl Tyrosine R =




Phosphoryl Serine R =




Phosphoryl Threonine R =

FIGURES 2A - 2C


2A



2B




2C




FIGURES 3A & 3B


3A



3B




FIGURES 4A - 4C



FIGURES 5A & 5B

5A



5B






FIGURE 6




FIGURES 7A - 7D


7A




7B



7C



7D



INTERNATIONAL SEARCH REPORT

In. national Application No

PCT/US 98/25196

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12Q1/42 C12Q1/42 G01N33/68 G01N33/573

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 98 18956 A (PANVERA CORP) 7 May 1998 see page 9, line 6, paragraph 2 - line 16 see example 4 see figure 2	1-35
A	WO 93 03377 A (ZILTENER HERMANN J) 18 February 1993 see examples 1,2	1-35
A	W J CHECOVICH, R E BOLGER, T BURKE: "Fluorescence polaritation - a new tool for cell and molecular biology" NATURE, vol. 375, 18 May 1995, pages 254-256, XP002100426 cited in the application see the whole document	1-35

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 April 1999

Date of mailing of the international search report

06/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Hart-Davis, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/25196

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Z-Y ZHANG, D MACLEAN, A M THIEME-SEFLER, R W ROESKE, J E DIXON: "A Continuous Spectrophotometric and Fluorimetric Assay for Protein Tyrosine Phosphatase Using Phosphotyrosine-Containing Peptides" ANALYTICAL BIOCHEMISTRY, vol. 211, 1993, pages 7-15, XP002100427 cited in the application see abstract; figures 1,3 ---	1-35
A	WO 95 18823 A (BETH ISRAEL HOSPITAL) 13 July 1995 see page 27; example 2 ---	1-35
A	WO 97 42501 A (BOEHRINGER INGELHEIM PHARMA) 13 November 1997 see examples 1-3 -----	1-35

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/25196

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9818956 A	07-05-1998	NONE	
WO 9303377 A	18-02-1993	NONE	
WO 9518823 A	13-07-1995	US 5532167 A	02-07-1996
WO 9742501 A	13-11-1997	CA 2253776 A EP 0901628 A	13-11-1997 17-03-1999